

SNUTP 02-017
 DESY 02-071
 hep-ph/0206049
 May 2002

Joint Angular Distribution of $B_s \rightarrow J/\psi\phi$ with Subsequent $\phi \rightarrow \rho\pi$ and $J/\psi \rightarrow \ell^+\ell^-$ Decays

G. Kramer

*II. Institut für Theoretische Physik, Universität Hamburg, D-22761 Hamburg,
 Germany*

H. S. Song

*Center for Theoretical Physics and School of Physics,
 Seoul National University, Seoul 151-742, Korea*

Chaehyun Yu

Department of Physics, Yonsei University, Seoul 120-749, Korea

The covariant density matrix formalism is used to obtain the joint angular distribution of the decay $B_s \rightarrow J/\psi\phi(1020)$ with subsequent $\phi \rightarrow \rho\pi$ and $J/\psi \rightarrow \ell^+\ell^-$ decays. The result is shown algebraically to be a special case of our previous work on the decay distribution of $B_d \rightarrow J/\psi K_1(1270)$ with $K_1 \rightarrow K\pi$ and $J/\psi \rightarrow \ell^+\ell^-$.

In the upgraded Tevatron Run II experiment, substantial numbers of $B_s \rightarrow J/\psi\phi(1020)$ events are expected in the future. Previous experimental works have been reported by the CDF Collaboration [1], and is well known that the decay process is described by three Lorentz invariant terms, *i.e.*, two CP-even terms and one CP-odd term. The joint angular distribution of $B_s \rightarrow J/\psi\phi$ with subsequent $\phi \rightarrow K^+K^-$ and $J/\psi \rightarrow \ell^+\ell^-$ decays can be described [2] in the same way as that of $B_d \rightarrow J/\psi K^*$ with subsequent $K^* \rightarrow$

$K\pi$ and $J/\psi \rightarrow \ell^+\ell^-$ decays [3], and a recent experimental investigation of the latter case has been reported by the Barbar [4] and the Belle [5] groups. Therefore, the joint angular distribution of $B_s \rightarrow J/\psi\phi$ will be determined with high precision in the future.

In the $B_s \rightarrow J/\psi\phi$ case, one expects $\phi \rightarrow \rho\pi$ and $\phi \rightarrow \pi\pi\pi$ in addition to the dominant $\phi \rightarrow K^+K^-$ decay. It will be difficult, if not impossible, to identify the $\phi \rightarrow \rho\pi$ events experimentally at Tevatron even though several hundred events are produced in the future. However, there is some possibility to construct a B_s factory in the future where the $\phi \rightarrow \rho\pi$ decay can be investigated more cleanly, and that would give additional information on the $B_s \rightarrow J/\psi\phi$ process.

Recently, the covariant density matrix formalism was used to obtain the joint angular distributions of $B_d \rightarrow J/\psi K^*(890)$, as well as $B_d \rightarrow J/\psi K_1(1270)$, followed by their subsequent decays [6]. The purpose of this brief report is to use this method to obtain the joint angular distribution of $B_s \rightarrow J/\psi\phi$ followed by $\phi \rightarrow \rho\pi$ and $J/\psi \rightarrow \ell^+\ell^-$ in the transversity basis [2, 3]. Since $B_d \rightarrow J/\psi K^*(890)$ and $B_s \rightarrow J/\psi\phi(1020)$ can be treated in the same way for $K^* \rightarrow K\pi$ and $\phi \rightarrow K^+K^-$, most of the results in Ref. 6 can be used to obtain the density matrix of J/ψ after the $\phi \rightarrow \rho\pi$ decay in the process $B_s \rightarrow J/\psi\phi(1020)$.

As shown in Ref. 6, the angular distribution of the decay $J/\psi \rightarrow \ell^+\ell^-$ is described in terms of the density matrix $\rho_{\lambda\lambda'}$ in the transversity basis, defined as in Fig. 1 in the J/ψ rest frame, as follows:

$$d\Gamma \sim \left[1 + \rho_{00} + (1 - 3\rho_{00}) \sin^2 \theta_{tr} \cos^2 \phi_{tr} \right. \\ + 2\text{Re}(\rho_{1-1})(\sin^2 \theta_{tr} \sin^2 \phi_{tr} - \cos^2 \theta_{tr}) \\ - 2\text{Im}(\rho_{1-1}) \sin 2\theta_{tr} \sin \phi_{tr} \\ + \sqrt{2}\text{Re}(\rho_{10} - \rho_{-10}) \sin^2 \theta_{tr} \sin 2\phi_{tr} \\ \left. - \sqrt{2}\text{Im}(\rho_{10} + \rho_{-10}) \sin 2\theta_{tr} \cos \phi_{tr} \right], \quad (1)$$

where θ_{tr} and ϕ_{tr} are the polar and azimuthal angles of the outgoing lepton ℓ^+ in the transversity basis as defined in Refs. 2 and 3, where the unit vectors $(\hat{n}_3, \hat{n}_1, \hat{n}_2)$ are along the $(\hat{x}, \hat{y}, \hat{z})$ axis.

The explicit values of the density matrix elements $\rho_{\lambda\lambda'}$ are calculated from the amplitude of the J/ψ production process. The matrix element of the $B_s \rightarrow J/\psi\phi$ decay is given in terms of three independent Lorentz scalars,

A , B and C , as [7, 8]

$$\mathcal{M} = A\epsilon_1^* \cdot \epsilon_2^* + \frac{B}{m_1 m_2} \epsilon_1^* \cdot k \epsilon_2^* \cdot q + \frac{iC}{m_1 m_2} \epsilon^{\mu\nu\sigma\tau} \epsilon_{1\mu}^* \epsilon_{2\nu}^* q_\sigma k_\tau, \quad (2)$$

where $\epsilon_i (i = 1, 2)$ are the polarization vectors of J/ψ and ϕ , respectively, and m_1, m_2 and q^μ, k^μ are their masses and momenta. Instead of scalar amplitudes A , B and C it is more convenient to introduce three transversity amplitudes $A_0, A_\parallel$ and $A_\perp$ to specify the CP property of the decay process. The latter are related to the former by [2]

$$\begin{aligned} A_0 &= -xA - (x^2 - 1)B, \\ A_\parallel &= \sqrt{2} A, \\ A_\perp &= \sqrt{2(x^2 - 1)} C, \end{aligned} \quad (3)$$

where x is defined as

$$x = \frac{k \cdot q}{m_1 m_2} = \frac{m_B^2 - m_1^2 - m_2^2}{2m_1 m_2}. \quad (4)$$

When the decay of the B_s into J/ψ and ϕ is specified by the polarization vectors $\epsilon_1(q, \lambda_1)$ and $\epsilon_2(k, \lambda_2)$, respectively, the joint density matrix elements of J/ψ and ϕ can be obtained from the square of Eq. (2) and by using the following relation [9] for each $\epsilon_i (i = 1, 2)$:

$$\epsilon^\mu(q, \lambda) \epsilon^{\nu*}(q, \lambda') = \frac{1}{3} \left(-g^{\mu\nu} + \frac{q^\mu q^\nu}{m_1^2} \right) \delta_{\lambda'\lambda} - \frac{i}{2m_1} \epsilon^{\mu\nu\sigma\tau} q_\sigma n_\tau^i (S^i)_{\lambda'\lambda} - \frac{1}{2} n_i^\mu n_j^\nu (S^{ij})_{\lambda'\lambda}, \quad (5)$$

where the (S^i) are the standard matrix representations of the spin-1 angular momentum operators and (S^{ij}) are traceless and symmetric matrices.

Then, one obtains

$$\begin{aligned} \rho_{\lambda_1 \lambda'_1, \lambda_2 \lambda'_2} &\sim \frac{1}{9} (|A_0|^2 + |A_\parallel|^2 + |A_\perp|^2) \delta_{\lambda_1 \lambda'_1, \lambda_2 \lambda'_2} \\ &\quad - \frac{2}{3\sqrt{\Delta}} \text{Re}(A_\parallel^* A_\perp) \{ m_1 n_1^i \cdot k (S^i)_{\lambda_1 \lambda'_1} \delta_{\lambda_2 \lambda'_2} + m_2 n_2^k \cdot q \delta_{\lambda_1 \lambda'_1} (S^k)_{\lambda_2 \lambda'_2} \} \\ &\quad + \frac{1}{\Delta} \{ (|A_\parallel|^2 + |A_\perp|^2) m_1 m_2 n_1^i \cdot k n_2^k \cdot q \end{aligned}$$

$$\begin{aligned}
& +\sqrt{\frac{\Delta}{2}}\text{Im}(A_0^*A_\perp)\langle qkn_1^in_2^k\rangle \\
& -\frac{\sqrt{2}}{4}\Delta\text{Re}(A_0^*A_\parallel)(n_1^i\cdot n_2^k - \frac{4q\cdot k}{\Delta}n_1^i\cdot kn_2^k\cdot q)\}(S^i)_{\lambda_1\lambda'_1}(S^k)_{\lambda_2\lambda'_2} \\
& -\frac{1}{3\Delta}(2|A_0|^2 - |A_\parallel|^2 - |A_\perp|^2)\{m_1^2n_1^i\cdot k n_1^j\cdot k (S^{ij})_{\lambda_1\lambda'_1}\delta_{\lambda_2\lambda'_2} \\
& \quad +m_2^2n_2^k\cdot q n_2^l\cdot q \delta_{\lambda_1\lambda'_1}(S^{kl})_{\lambda_2\lambda'_2}\} \\
& -\frac{1}{\Delta}\{ \sqrt{2}\text{Im}(A_0^*A_\parallel)\langle qkn_1^in_2^k\rangle \\
& \quad +\frac{2}{\sqrt{\Delta}}\text{Re}(A_\parallel^*A_\perp)m_1m_2n_1^i\cdot kn_2^k\cdot q \\
& \quad -\sqrt{\frac{\Delta}{2}}\text{Re}(A_0^*A_\perp)(n_1^i\cdot n_2^k - \frac{4q\cdot k}{\Delta}n_1^i\cdot kn_2^k\cdot q)\} \\
& \quad \times\{m_2n_2^l\cdot q (S^i)_{\lambda_1\lambda'_1}(S^{kl})_{\lambda_2\lambda'_2} + m_1n_1^j\cdot q (S^{ij})_{\lambda_1\lambda'_1}(S^k)_{\lambda_2\lambda'_2}\} \\
& +\{ \frac{4}{\Delta^2}|A_0|^2m_1^2m_2^2n_1^i\cdot kn_1^j\cdot kn_2^k\cdot qn_2^l\cdot q \\
& \quad +\frac{1}{8}|A_\parallel|^2(n_1^i\cdot n_2^k - \frac{4q\cdot k}{\Delta}n_1^i\cdot kn_2^k\cdot q)(n_1^j\cdot n_2^l - \frac{4q\cdot k}{\Delta}n_1^j\cdot kn_2^l\cdot q) \\
& \quad +\frac{1}{2\Delta}|A_\perp|^2\langle qkn_1^in_2^k\rangle\langle qkn_1^jn_2^l\rangle \\
& \quad -\frac{\sqrt{2}}{2\Delta}\text{Re}(A_0^*A_\parallel)m_1m_2n_1^i\cdot kn_2^k\cdot q(n_1^j\cdot n_2^l - \frac{4q\cdot k}{\Delta}n_1^j\cdot kn_2^l\cdot q) \\
& \quad +\frac{2\sqrt{2}}{\Delta\sqrt{\Delta}}\text{Im}(A_0^*A_\perp)m_1m_2n_1^i\cdot kn_2^k\cdot q\langle qkn_1^jn_2^l\rangle \\
& \quad -\frac{1}{2\sqrt{\Delta}}\text{Im}(A_\parallel^*A_\perp)(n_1^i\cdot n_2^k - \frac{4q\cdot k}{\Delta}n_1^i\cdot kn_2^k\cdot q)\langle qkn_1^jn_2^l\rangle\} \\
& \quad \times(S^{ij})_{\lambda_1\lambda'_1}(S^{kl})_{\lambda_2\lambda'_2}. \tag{6}
\end{aligned}$$

Here, the density matrix is multiplied by $(|A_0|^2 + |A_\parallel|^2 + |A_\perp|^2)$ for convenience, and Δ and $\langle qkn_1^jn_2^l\rangle$ are defined as

$$\begin{aligned}
\Delta &= (m_B^4 + m_1^4 + m_2^4 - 2m_B^2m_1^2 - 2m_B^2m_2^2 - 2m_1^2m_2^2) \\
&= 4[(q\cdot k)^2 - m_1^2m_2^2] = 4m_1^2m_2^2(x^2 - 1), \tag{7}
\end{aligned}$$

$$\langle qkn_1^jn_2^l\rangle = \epsilon^{\mu\nu\sigma\tau}q_\mu k_\nu(n_\sigma^j)_1(n_\tau^l)_2. \tag{8}$$

The amplitude of the decay process $\phi(k, \lambda_2) \rightarrow \rho(\tilde{k}')\pi(\tilde{k})$ is described by

$$\mathcal{M}_2 = if\epsilon^{\mu\nu\lambda\tau}\epsilon_{2\mu}(k)\epsilon_{3\nu}^*(\tilde{k}')k_\lambda\tilde{k}_\tau, \quad (9)$$

where ϵ_3^μ and $\tilde{k}'$ are the polarization vector and the momentum of ρ , respectively. After the polarization of ρ is summed over, one obtains

$$\rho_{\lambda'_2\lambda_2}^D(\phi) = \frac{1}{3}\delta_{\lambda'_2\lambda_2} + \frac{m_\phi^2}{\bar{\Delta}}\tilde{k}\cdot n_i\tilde{k}\cdot n_j(S^{ij})_{\lambda'_2\lambda_2}, \quad (10)$$

where $\bar{\Delta}$ is defined as

$$\bar{\Delta} = m_\phi^4 + m_\rho^4 + m_\pi^4 - 2m_\phi^2 - 2m_\rho^2 - 2m_\pi^2. \quad (11)$$

The result of Eq. (10) can be obtained from Eq. (6). Then, the density matrix of J/ψ can be obtained from the product

$$\rho_{\lambda_1\lambda'_1,\lambda_2\lambda'_2}\rho_{\lambda'_2\lambda_2}^D(\phi). \quad (12)$$

The joint angular distribution for the decay process $B_s \rightarrow J/\psi\phi$ ($J/\psi \rightarrow \ell^+\ell^-$, $\phi \rightarrow \rho\pi$) in the transversity basis becomes

$$\begin{aligned} \frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta_1 d\cos\theta_{tr} d\phi_{tr}} &= \frac{9}{64\pi} \left[2|A_0|^2 \left\{ \sin^2\theta_1(1 - \sin^2\theta_{tr}\cos^2\phi_{tr}) \right\} \right. \\ &\quad + |A_{\parallel}|^2 \left\{ \cos^2\theta_1(1 - \sin^2\theta_{tr}\sin^2\phi_{tr}) + \sin^2\theta_{tr} \right\} \\ &\quad + |A_{\perp}|^2 \left\{ 1 - \sin^2\theta_{tr}\sin^2\phi_{tr} + \cos^2\theta_1\sin^2\theta_{tr} \right\} \\ &\quad - \text{Im}(A_{\parallel}^*A_{\perp}) \sin^2\theta_1 \sin 2\theta_{tr} \sin\phi_{tr} \\ &\quad + \frac{1}{\sqrt{2}}\text{Re}(A_0^*A_{\parallel}) \sin 2\theta_1 \sin^2\theta_{tr} \sin 2\phi_{tr} \\ &\quad \left. - \frac{1}{\sqrt{2}}\text{Im}(A_0^*A_{\perp}) \sin 2\theta_1 \sin 2\theta_{tr} \cos\phi_{tr} \right], \quad (13) \end{aligned}$$

where θ_1 is the angle between the ϕ direction and the π direction in the ϕ rest frame, and the three amplitudes are normalized as $|A_0|^2 + |A_{\parallel}|^2 + |A_{\perp}|^2 = 1$. It is noted that in Ref. 6 we have considered the process $B \rightarrow J/\psi K_1(1270)$ and $J/\psi \rightarrow \ell^+\ell^-$, $K_1 \rightarrow K\pi$ and Eq. (13) is a special case obtained by putting $\alpha = -1$ in Eq. (28) or $\xi = 3/2$ in Eq. (29) of Ref. 6. This method can be used to discuss the polarization effects in W boson productions at e^+e^- linear collider [10].

Acknowledgments


The authors would like to thank Prof. Intae Yu for discussion related with the CDF experiments. The work is supported in part by the DFG-KOSEF Collaboration (2000) under contracts 446 KOR-113/137/0-1 (DFG) and 20005-111-02-2 (KOSEF). HS was supported in part by the Korea Research Foundation (KRF-2000-D00077) and by the BK21 program. The work of CY was supported by grant No. R02-2002-000-00168-0 of the KRF. The authors thank their guest institutes for the warm hospitality during their visits.

References

- [1] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **75**, 1451 (1995); Phys. Rev. D **53**, 3496 (1996).
- [2] I. Dunietz, H. J. Lipkin, H. Quinn, A. Snider, and W. Toki, Phys. Rev. D **43**, 2193 (1991).
- [3] A. S. Dighe, I. Dunietz, H. J. Lipkin, and J. L. Rosner, Phys. Lett. B **369**, 144 (1996).
- [4] Babar Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 241801 (2001).
- [5] Belle Collaboration, K. Abe *et al.*, BELLE-CONF-0105 (2001).
- [6] G. Kramer, H. S. Song, and Chaehyun Yu, Phys. Rev. D **65**, 091501(R) (2002); *ibid.* **66**, 095901(E) (2002).
- [7] G. Kramer and W. F. Palmer, Phys. Rev. D **45**, 193 (1992).
- [8] G. Valencia, Phys. Rev. D **39**, 3339 (1989).
- [9] S. Y. Choi, Taeyeon Lee, and H. S. Song, Phys. Rev. D **40**, 2477 (1989).
- [10] D. W. Jung, K. Y. Lee, H. S. Song, and C. Yu, J. Korean Phys. Soc. **36**, 258 (2000); Czech. J. Phys. **51**, A173 (2001).

Figure Captions

Fig. 1. Definitions of the angles in the transversity basis. θ_{tr} and ϕ_{tr} are the polar and the azimuthal angles of ℓ^+ in the J/ψ rest frame. θ_1 is the angle between the ϕ direction and the π direction in the ϕ rest frame.



8